

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

calcd 301.2406, found 301.2402.

RBL-1 5-Lipoxygenase Inhibition Assay. 5-Lipoxygenase activity was measured in the 20000g supernatant from homogenized rat basophilic leukemia (RBL-1) cells. Inhibitors or vehicle (2% Me₂SO) were preincubated for 20 min with the RBL-1 supernatant (7.5 × 10⁶ cell equivalents/mL) at 37 °C in pH 6.8 buffer (10 mM BES, 10 mM PIPES, 1 mM EDTA, 0.1 M NaCl, 0.7 mM CaCl₂) prior to initiating the 5-lipoxygenase reaction by addition of 66 μM [¹⁴C]arachidonic acid. [³H]-5-HETE added to the reaction mixture served as a recovery standard. Reactions were terminated by acidification to pH 3 and the mixtures were extracted with diethyl ether. The ether extracts were evaporated under nitrogen and the reaction products were separated from nonconverted substrate by thin-layer chromatography. Radioactivity comigrating with 5-HETE was measured by liquid scin-

tillation counting and corrected for recovery of [³H]-5-HETE. Inhibition was calculated as the percent reduction from control levels of [¹⁴C]-5-HETE formation. Concentrations causing 50% inhibition (IC₅₀'s) and their 96% confidence limits were calculated as the 50% intercept and their fiducial limits from linear regression analysis²² of percent inhibition vs. log concentration plots.

Acknowledgment. We thank Dr. R. Walters for helpful discussions, D. Bornemeier for excellent technical assistance, and the spectroscopic services department at Abbott for the NMR and MS data.

(22) Ostle, B. *Statistics in Research*, 2nd ed., The Iowa State University Press: Ames, IA, 1963; pp 159-221.


Structure-Activity Relationships among Di- and Tetramine Disulfides Related to Benextramine

M. Alvarez,* R. Granados, D. Mauleón, G. Rosell, M. Salas, J. Sallés, and N. Valls

Departamento de Química Orgánica y Farmacéutica, Facultad de Farmacia, Universidad de Barcelona, Barcelona 08028, Spain. Received January 10, 1986


The synthesis and irreversible α-blocking activity in the rat vas deferens of a series of tetra- and diamine disulfides 2-38, structural analogues of benextramine (BHC), are described. All compounds containing a central cystamine moiety displayed an irreversible α-adrenergic blockade at concentrations ranging from 10⁻⁴ to 6 × 10⁻⁶ M. Potency was increased in cystamines N,N'-disubstituted with 6-aminoethyl groups, especially when the outer nitrogen atoms bear arylalkyl substituents or are enclosed in a ring. However, N,N,N',N'-tetrasubstituted cystamines were poor blockers. Structural specificity in the outer portion of the tetramine disulfide is low, since many types of substituents gave rise to potent α-blockers. Even replacement of the outer amines with nonbasic ethers or amides was observed to maintain irreversible α-blockade.

In a series of structure-activity relationship (SAR) studies of tetramine disulfides carried out by Melchiorre and Belleau,¹ optimum α-adrenergic blocking potency was found for compounds N,N'-bis[6-[(o-methoxybenzyl)-amino]hexyl]cystamine (BHC, 1) and N,N'-bis(8-amino-octyl)cystamine, (AOC, 39). From the structural features of these compounds, the authors proposed a topographical model for the α-adrenoceptor in the rat vas deferens.



Accordingly, BHC would interact with a set of four anionic centers and two flat areas complementary to the aromatic rings, and AOC would bind to a different set of four anionic centers. These two binding areas would be symmetrically disposed on the surface of the receptor and share a common central thiol group. According to the hypothesis, the initial electrostatic interactions between the four cationic nitrogen atoms and the anionic centers would lead to a conformational change unmasking the thiol

group. This would allow a disulfide-thiol exchange reaction, which in turn results in a covalent blockade of the α-adrenoceptor. The analogues of tetramine disulfides BHC and AOC showed different SAR. Thus, the optimum chain length between the "inner" and "outer" nitrogen atoms was found to be six carbon atoms in BHC and eight methylene groups in AOC. Substitution on the outer nitrogen atom led to a reduced potency in AOC, while in BHC a benzylic substituent (especially an o-methoxybenzyl group) on this position gave maximal potency. On the other hand, methylation of the inner nitrogen atoms led to a marked reduction in potency in the BHC series while the AOC methylated analogues showed almost no changes in activity. On the basis of the high α-blocking activity of the catechol-containing disulfide 40, which was




equiactive with BHC, Melchiorre also suggested a common binding site for the outer (3,4-dihydroxybenzyl)amino moiety of 40, as well as the (o-methoxybenzyl)amino group of BHC and the catecholamine neurotransmitters, in spite of the benzylic nature of the N-substituent and the lack of a benzylic hydroxyl group found in catecholamines.

Aiming to extend and evaluate the above possibilities and especially the predicted relationship between the catecholamine and BHC binding sites, we undertook the synthesis of a new series of polyamine disulfides containing terminal moieties structurally related in some instances to α-adrenergic drugs.² Thus, compounds 2-9 (Table I)

- (1) (a) Melchiorre, C.; Yong, M. S.; Benfey, B. G.; Belleau, B. *J. Med. Chem.* 1978, 21, 1126. (b) Melchiorre, C.; Giardina, D.; Brasili, L.; Belleau, B. *Farmacol.* 1978, 33, 999. (c) Melchiorre, C.; Giannella, M.; Brasili, L.; Benfey, B. G.; Belleau, B. *Eur. J. Med. Chem.* 1981, 16, 111. (d) Melchiorre, C. *Trends Pharmacol. Sci.* 1981, 2, 209. (e) Angeli, P.; Brasili, L.; Brancia, E.; Giardina, D.; Quaglia, W.; Melchiorre, C. *J. Med. Chem.* 1985, 28, 1643. (f) Bertini, R.; Giardina, D.; Gullini, U.; Pignini, M.; Melchiorre, C.; Carpy, A. *Eur. J. Med. Chem.* 1985, 20, 309.

Scheme I




are structural combinations of BHC and phenylethanolamines; in disulfide 9 the group replacing the (o-methoxybenzyl)amino substituent is specifically norepinephrine. Compound 11 is related to amphetamine and disulfides 10 contain the α -blocking (1,4-benzodioxan-2-ylmethyl)-amino moiety.

The results of pharmacological testing of the above disulfides prompted us to carry out our synthesis of new BHC analogues bearing modified outer portions. The changes made were related to (a) change in the substitution pattern of the aromatic ring and in the distance between this ring and the outer nitrogen atom (14-16), (b) replacement of heterocycles for the aryl group (17-20), (c) change of the amino groups into nonbasic amide or ether functions (12, 13, 30-34), (d) replacement of nonaromatic heterocycles for the benzylamino moiety (21-24), (e) introduction of a new ring linking the Ar/N moieties (25-29), and (f) suppression of the whole outer moiety, leading to N-substituted cystamines (35-38).

Chemistry

Preparation of symmetrical disulfides 2-10, 12-29 (see specific structures in Table I), and 49^{1c} is outlined in Scheme I. In method A, reaction of 3-(6-aminohexyl)-1,3-thiazolidine (42)^{2a} with an appropriate alkylating agent (epoxides 41a-h^{2a} for 2-9 and racemic^{2a} or optically pure^{2b} tosylate 43 for 10) gave the required thiazolidines 45

Scheme II. Method D



(structures in Table II). Oxidative ring opening and dimerization by treatment with 0.1 N I₂/KI² yielded disulfides 2-10. Compounds 12 and 13 were obtained from (aminoalkyl)thiazolidine 42 by acylation with acyl chlorides 44a,c and oxidative coupling. Reduction of amide group of 12 with LiAlH₄ gave the cystamine 14. Simultaneous reduction of amide and nitro groups of the cystamine obtained from thiazolidine 45k gave 15. Ring opening and reduction of the amide group of thiazolidine 45l with LiAlH₄ gave a (methylamino)ethanethiol, which upon reaction with I₂ gave the N-methylated hexamine 16. The tetrahydroquinazoline derivatives 25 and 26 and the benzoxazine 27 were obtained by ring closure of o-amino-benzylamines 15 and 16 or hydroxybenzylamine 49, respectively, with HCHO in alkaline solution.




The heteroarylmethyl-substituted disulfides 17-20 were obtained from N,N'-bis(6-aminohexyl)cystamine (46)^{1a} through method B consisting of a reductive alkylation of the primary amino groups with an appropriate heterocyclic aldehyde, 47.

In method C the starting materials were the terminal heterocycle (piperidine, pyrrolidine, tetrahydroisoquinoline, or indoline) and 6-phthalimidohexanoyl chlo-

- (2) The preparation of some of these compounds has been previously described; see: (a) Granados, R.; Alvarez, M.; Valls, N.; Sales, M. *J. Heterocycl. Chem.* 1983, 20, 1271. (b) Granados, R.; Valls, N.; Alvarez, M.; Bardaji, E. *An. Quim., Ser. C* 1983, 79, 303. (c) Alvarez, M.; Granados, R.; Lavilla, R.; Sales, M. *J. Heterocycl. Chem.* 1983, 22, 145. (d) Alvarez, M.; Granados, R.; Rosell, G.; Santaló, P.; Sales, M. *An. Quim., Ser. C* 1984, 80, 283.
- (3) (a) Koo, J.; Avacain, S.; Martin, G. *J. Am. Chem. Soc.* 1965, 77, 5373. (b) Nelson, W. L.; Wennerstrom, J. E.; Dyer, D. C.; Engel, U. *J. Med. Chem.* 1977, 20, 880.

Table I. Structures of the Tetra- and Diamine Disulfides and α -Adrenergic Blocking Results

no.	X	Y	Z	W	n	yield, % (method)	mp, °C	purifn solvent	formula ^a	α -blockade ^b		
										10 ⁻⁴ M	2 × 10 ⁻⁴ M	6 × 10 ⁻⁶ M
1/						ref 1a						
2	H	H	H	H	H	75 (A)	180-181	EtOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ·3H ₂ O	40	100	35
3	H	-OCH ₃ -	H	H	H	75 (A)	155-160	MeOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·3C ₂ H ₅ O ₄ ·2H ₂ O	13	43	
4	OMe	H	H	H	H	88 (A)	133-136	MeOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ·2H ₂ O	82	91	
5	H	OMe	H	H	H	77 (A)	135-138	MeOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·3C ₂ H ₅ O ₄ ·H ₂ O	91	81	
6	H	OMe	OMe	H	H	54 (A)	135-138	MeOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄	87	98	
7	H	OMe	OMe	H	OMe	88 (A)	200-203	MeOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄	87	98	
8	H	OMe	OMe	OMe	OMe	89 (A)	200-203	MeOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ·H ₂ O	92		
9	H	OH	OH	H	H	60 (A)	107-110	EtOH	C ₂₃ H ₃₄ N ₄ O ₈ S ₂ ·4HCl·C ₂ H ₅ O ₄			
10	(1,4-benzodioxan-2-yl- methyl)amino	H	H	H	6	61 (A)	260-262	Et ₂ O	C ₃₄ H ₄₄ N ₄ O ₈ S ₂ ·4HCl	53	inact	
(S,S)-10	(1,4-benzodioxan-2-yl- methyl)amino	H	H	H	6	63 (A)	262-264	MeOH-Et ₂ O	C ₃₄ H ₄₄ N ₄ O ₈ S ₂ ·4HCl	79	inact	
(R,R)-10	(1,4-benzodioxan-2-yl- methyl)amino	H	H	H	6	65 (A)	262-264	MeOH-Et ₂ O	C ₃₄ H ₄₄ N ₄ O ₈ S ₂ ·4HCl	68	inact	
11	C ₆ H ₅ CH ₂ CH(NH ₂)	H	H	H	5	56 (D)	152-155	MeOH	C ₃₀ H ₄₀ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄		57	
12	C ₆ H ₅ CH ₂ CONH	H	H	H	6	51 (A)	184-188	MeOH	C ₃₇ H ₅₀ N ₄ O ₈ S ₂ ·2C ₂ H ₅ O ₄ ·CH ₃ OH		44	
13	o-NH ₂ C ₆ H ₄ CONH	H	H	H	6	37 (A)	indef	EtOH	C ₃₀ H ₄₀ N ₄ O ₈ S ₂ ·4HCl ^f	55	inact	
14	C ₆ H ₅ CH ₂ CH ₂ NH	H	H	H	6	95 (A)	194-196	MeOH	C ₃₂ H ₄₄ N ₄ O ₈ S ₂ ·3C ₂ H ₅ O ₄ ·H ₂ O	32	inact	
15	o-NH ₂ C ₆ H ₄ CH ₂ NH	H	H	H	6	43 (A)	134-136	MeOH	C ₃₀ H ₄₀ N ₄ O ₈ S ₂ ·6HCl·C ₂ H ₅ OH	79	inact	
16	o-NH ₂ C ₆ H ₄ CH ₂ NH	Me	H	H	6	56 (A)	210-215	EtOH	C ₃₂ H ₄₄ N ₄ O ₈ S ₂ ·6HCl	38	inact	
17	2-phenylamino	H	H	H	6	67 (B)	275-277	H ₂ O	C ₂₅ H ₃₄ N ₄ O ₈ S ₂ ·4HCl	70	inact	
18	furfurylamino	H	H	H	6	75 (B)	221-223	MeOH-H ₂ O	C ₂₅ H ₃₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄	100		79
19	3-indolylmethylamino	H	H	H	6	80 (B)	186-190	EtOH	C ₃₄ H ₄₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄		inact ^d	
20	[(1-methyl-3-pyrrolyl)- methyl]amino	H	H	H	6	80 (B)	indef	MeOH	C ₃₄ H ₄₄ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ^f	46 ^d	100	54
21	piperidino	H	H	H	6	50 (C)	121-123	MeOH	C ₂₈ H ₄₀ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ·H ₂ O		100	56
22	1-pyrrolidinyl	H	H	H	6	45 (C)	120-121	MeOH	C ₂₈ H ₄₀ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄		96	
23	1-piperidinyl	Me	Me	Me	6	54 (C)	229-231	EtOH	C ₂₈ H ₄₀ N ₄ O ₈ S ₂ ·4HCl·2H ₂ O	34	41	
24	1-pyrrolidinyl	Me	Me	Me	6	50 (C)	185-188	EtOH	C ₂₈ H ₄₀ N ₄ O ₈ S ₂ ·4HCl·2H ₂ O	34	41	
25	NH	H	H	H	1	80 (A)	69-70	EtOH	C ₂₇ H ₃₈ N ₄ O ₈ S ₂ ·6C ₂ H ₅ N ₃ O ₇	89	inact	
26	NH	Me	Me	Me	1	84 (A)	80-85	MeOH	C ₂₈ H ₄₀ N ₄ O ₈ S ₂ ·6C ₂ H ₅ O ₄	6	inact ^d	
27	O	H	H	H	1	78 (B)	164-167	MeOH	C ₂₇ H ₃₈ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ·CH ₃ OH	34 ^d	inact ^d	
28	CH ₃	H	H	H	1	40 (C)	105-108	MeOH	C ₂₈ H ₄₀ N ₄ O ₈ S ₂ ·4C ₂ H ₅ O ₄ ·3H ₂ O	66 ^d	inact ^d	




29	CH ₃	H	35 (C)	158-160	MeOH	C ₂₇ H ₃₀ N ₂ S ₂ ·2C ₂ H ₅ O ₄	37
30	C ₂ H ₅ CH ₃	NH	15 (E)	192-194	MeOH	C ₃₀ H ₄₄ N ₂ O ₇ ·2C ₂ H ₅ O ₄	32
31	o-MeOC ₂ H ₄ CH ₃	NH	72 (E)	138-140	MeOH	C ₂₃ H ₃₂ N ₂ O ₅ ·S ₂ ·2C ₂ H ₅ O ₄	36
32	C ₂ H ₅ CH ₃	O	60 (F)	201-204	Me·CO	C ₃₂ H ₄₄ N ₂ O ₅ ·S ₂ ·2HCl	60
33	o-MeOC ₂ H ₄ CH ₃	O	80 (F)	198-200	MeOH	C ₂₃ H ₃₂ N ₂ O ₅ ·S ₂ ·2C ₂ H ₅ O ₄	58
34	3,4-(methylethenioxy)-benzyl	O	61 (F)	185-188	MeOH-H ₂ O	C ₂₂ H ₃₀ N ₂ O ₅ ·S ₂ ·2C ₂ H ₅ O ₄	45
35	C ₂ H ₅	Et	48 (G)	88-90	EtOH	C ₁₉ H ₂₈ N ₂ S ₂ ·2HCl	inact ^a
36	t-Bu	H	10 (G)	197-199	EtOH	C ₁₉ H ₂₈ N ₂ S ₂ ·2HCl	inact ^a
37	C ₂ H ₅ CH ₃	H	55 (G)	263-265	MeOH-H ₂ O	C ₁₉ H ₂₄ N ₂ S ₂ ·2HCl	48
38	-(CH ₂) ₇ -		83 (G)	249-251	MeOH	C ₁₄ H ₂₀ N ₂ S ₂ ·2HCl	inact ^a

* All compounds were analyzed for C, H, and N and analytical values were within $\pm 0.4\%$ of calculated values. ^b Potencies as irreversible inhibitors of NE-induced response in rat vas deferens with BHC as standard (see Experimental Section). All compounds were tested at least five times and the percent inhibition is accurate to $\pm 5\%$. ^c These compounds have indefinite melting points. ^d Precipitation in the organ bath was observed during the incubation period. ^e These disulfides were tested also at a 10^{-3} M concentration, showing the blockade values of 34% (35), 84% (36), and 83% (38). ^f Benextramine (BHC).

Scheme III

method E
(for compounds 20, 21)

method F
(for compounds 82-84)



ride. The phthalimido protecting group in the resulting amides was removed by hydrazine treatment to give the amino amides 48a-d. Reduction of amino amides 48a-d with LiAlH_4 followed by condensation with thiirane and oxidative dimerization yielded compounds 21, 22, 28, and 29. The *N*-methyl derivatives 23 and 24 were also obtained from 48a,b through *N*-formylation with AcOCHO , reduction of resulting formamides 48e,f with LiAlH_4 , and the subsequent steps of method C.

The amphetamine-related tetramine disulfide 11 was synthesized following method D (Scheme II). Alkylation of 2-benzyl-1,3-dithiane⁴ with 1,5-dibromopentane and elimination of the protective group afforded a bromo ketone. This compound was used for alkylation of 1,3-azolidine to give the compound 50. Oxidative ring opening of 50 by I_2/KI treatment gave the dioxo disulfide 51. The carbonyl groups in disulfide 51 were easily transformed into amino groups by their conversion to oximes and reduction with $LiAlH_4$.

Disulfides 30 and 31, whose inner nitrogen atoms have been replaced by oxygen atoms, were prepared as indicated in Scheme III (method E). The starting material, S-protected mercaptoethanol⁵ 52, was O-alkylated with 1,5-dibromopentane. The remaining bromine atom was displaced by a cyano group that served as the precursor of the primary amino function in 53. Cleavage of the C-S benzylic bond with Na in EtOH followed by oxidative coupling of the resulting thiol yielded disulfide 54 having the required functionalization. The last step of method E was the introduction of the benzylic N-substituents achieved through reductive alkylation of disulfide 54 with benzaldehyde or o-methoxybenzaldehyde.


Amino ethers 32-34, lacking the external nitrogen atoms, were obtained through method F (Scheme III) with alcohols 55a-c as the starting material, respectively. Alkylation of the corresponding alkoxide with 1,6-dibromohexane, followed by reaction with potassium phthalimide and removal of the *N*-phthaloyl group, yielded the amino ethers 56. These compounds were converted into 32-34 by mercaptoethylation with thiirane and oxidative dimerization with $K_2Fe(CN)_6$.

Finally, cystamines 35–38 were synthesized in two steps (method G) from the corresponding amine (diethylamine,

(4) Nagao, Y.; Seno, K.; Fujita, E. *Tetrahedron Lett.* 1979, 4403.

(5) Szabo, J. L.; Sciller, R. T. *J. Am. Chem. Soc.* 1948, 70, 3667.

Table II. Structures and Physical Properties of Compounds 45, 48, 50, 51, 53, 54, and 56



no.	X	Y	Z	W	n	yield, %	mp, °C	purifn solvent	formula ^a
45a	H	H	H	H		81	129-135	EtOH	C ₁₇ H ₂₉ N ₂ O ₅ S-2C ₂ H ₅ O ₄ ·3H ₂ O
45b	H		-OCH ₂ O-	H		49	93-95	MeOH-Et ₂ O	C ₁₈ H ₂₉ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45c	OMe	H	H	H		81	75-80	MeOH-Et ₂ O	C ₁₈ H ₂₉ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45d	H	OMe	H	H		62	50-52	MeOH-Et ₂ O	C ₁₉ H ₃₁ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45e	H	H	OMe	H		64	134-136	MeOH-Et ₂ O	C ₁₉ H ₃₁ N ₂ O ₅ S-2C ₂ H ₅ O ₄ ·C ₄ H ₁₀ O
45f	H	OMe	OMe	H		66	162-165	MeOH-Et ₂ O	C ₁₉ H ₃₁ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45g	H	OMe	OMe	OMe		77	100-102	MeOH-Et ₂ O	C ₁₉ H ₃₁ N ₂ O ₅ S-2C ₂ H ₅ O ₄ ·CH ₃ OH
45h	H	OCH ₂ OCH ₃	OCH ₂ OCH ₃	H		75	68-70	MeOH-Et ₂ O	C ₂₀ H ₃₃ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45i						86	198-200	MeOH	C ₂₁ H ₃₅ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45j	H				1	92	144-146	MeOH	C ₁₈ H ₂₉ N ₂ O ₅ S-2C ₂ H ₅ O ₄
45k	NO ₂				0	99	173-175	MeCN	C ₁₇ H ₂₇ N ₂ O ₅ S-C ₂ H ₅ O ₄ ·CH ₃ OH
45l	NH ₂				0	92	indef	EtOH	C ₁₈ H ₂₉ N ₂ O ₅ S-HCl
48a						72	95-97	MeOH-Me ₂ CO	C ₁₈ H ₂₉ N ₂ O ₅ S-HCl ^b
48b						70	121-122	MeOH-Et ₂ O	C ₁₁ H ₂₁ N ₂ O ₅ C ₂ H ₅ O ₄
48c						70	110-115	MeOH	C ₁₀ H ₂₀ N ₂ O ₅ C ₂ H ₅ O ₄
48d						60	200-215	MeOH-H ₂ O	C ₁₅ H ₂₅ N ₂ O ₅ C ₂ H ₅ O ₄ ·CH ₃ OH
48e						69	200 ^c		C ₁₄ H ₂₃ N ₂ O ₅ C ₂ H ₅ O ₄
48f						61	190 ^c		C ₁₃ H ₂₁ N ₂ O ₅
50						41	145-149	MeOH	C ₁₁ H ₂₁ N ₂ O ₅
51						97	192-195	MeOH	C ₁₆ H ₂₇ NOS-C ₂ H ₅ O ₄
53						14	97-100	MeOH	C ₂₄ H ₄₉ N ₂ O ₅ S-2C ₂ H ₅ O ₄
54						90	d		C ₁₆ H ₂₇ NOS-C ₂ H ₅ O ₄ ·CH ₃ OH
56a						85	118-120	Me ₂ CO	C ₁₆ H ₂₇ N ₂ O ₅ S ₂
56b						85	110-112	Me ₂ CO	C ₁₃ H ₂₁ NO-C ₂ H ₅ O ₄
56c						85	175-180	Me ₂ CO	C ₁₄ H ₂₃ NO ₂ -C ₂ H ₅ O ₄

^a All compounds were analyzed for C, H, and N and analytical values were within $\pm 0.4\%$ of calculated values. ^b This compound has an indefinite melting point. ^c Boiling point at 0.07 mmHg. ^d Purified by column chromatography.

tert-butylamine, benzylamine, or piperidine) by mercaptoethylation with thirane and oxidative coupling of the resulting thiols by I₂/KI treatment.

Results and Discussion

The disulfides 1-38 were tested as irreversible blockers of the norepinephrine-induced contractions in rat as deferens. With benextramine (BHC, 1) as a standard for comparison, the relative potencies observed are assembled in Table I.

A blockade of 91% of the NE contraction, almost equivalent to the inhibition caused by BHC, was found for the catechol-containing disulfide 9, a molecular combination of BHC and NE. This result does not disagree with the hypothesis of Melchiorre attributing a single binding site for the catecholamines and the outer moieties of tetramine disulfides. Although it is well-known that methylation of the phenolic groups of catecholamines greatly reduces their α -adrenergic affinity, this result was not found in our series of phenylethanolamine-bearing disulfides. Even though the aromatic ring of compounds 4-8 contains one, two, or three methoxy groups and no hydroxyl functionalities, these disulfides are potent blockers, especially the trimethoxy derivative, which almost completely inhibits the NE-induced response. Compounds 4-8 could be seen as structural analogues of some methoxyl-containing α -blockers, such as WB-4101. However, their binding is not very specific since the number and/or position of methoxyl groups does not greatly influence their adrenergic blocking potency. On the other hand, Melchiorre has recently reported¹¹ a study on the molecular combination of WB-4101 and BHC and concluded that these two kinds of α -blockers have different binding sites.

In good agreement with the above results, 1,4-benzodioxan-2-ylmethyl-substituted tetramine disulfide 10 showed a very poor blocking activity. The stereoselectivity of α -adrenoceptors toward competitive antagonists such as 2-(aminocethyl)-1,4-benzodioxanes is a well-established phenomenon.^{3b} In these drugs, 2*S* isomers are consistently more potent than the corresponding 2*R* enantiomers. The adrenergic blocking potency of the enantiomeric benzodioxane-containing disulfides (*S,S*)-10 and (*R,R*)-10 was found to be very low at 2×10^{-6} M and almost equivalent for both isomers at higher concentrations. The low activity of derivatives 10 suggests that either the normal benzodioxan blocking site is not being occupied or that the molecule cannot orient itself properly to take advantage of optimum binding at the site. The small difference in activity between optical isomers (*S,S*)-10 and (*R,R*)-10 also suggests that the binding is not very specific.

Isosteric replacement of the *o*-methoxyl groups in BHC for amino groups, as in 15, or formation of a new ring between the outer nitrogen atoms and the aromatic ring, as in 25, did not bring about a significant change in blocking activity. Again, these results suggest that the flexibility of the binding site can accommodate structural changes such as substitution of nitrogen for oxygen or ring closed systems. The benzoxazinyl and isoquinolyl derivatives 27 and 28 were only slightly soluble in the Krebs solution and therefore did not give reliable results.

Further evidence for the ability of the tetramine disulfide binding site to accommodate many types of compounds is given by the high blocking activity of some heteroarylmethyl-substituted tetramines. Furan and pyrrole analogues 18 and 20 show even greater potency than BHC at 6×10^{-6} M. The low blockade exerted by

19 can be ascribed again to lack of solubility since we observed a precipitate in the organ bath during the incubation period. Although the heteroarylmethyl moieties of compounds 17–20 do not bear any structural resemblance to the adrenergic aryloethanolamines, they could bind strongly with a site capable of accommodating flat aromatic substituents, similar to those found in BHC and in conventional α -antagonists. However, this terminal site has very low structural specificity since it can bind efficiently to compounds possessing a saturated heterocyclic ring, as in piperidinyl and pyrrolidinyl derivatives 21 and 22. Lacking an aromatic rings does not reduce their blocking potencies, given that 21 is even more potent than BHC. On the other hand, 1-indolinyl-substituted disulfide 29 is only a mild blocker even though it has an aromatic ring condensed to the heterocycle and thus is structurally more similar to BHC than 22. It could be thought that disulfides 21–24 are not related to BHC, but to (amino-octyl)cystamine (AOC), also a potent blocker in the rat vas deferens. Nevertheless, 21 and 22 retain the structure-activity relationships described for BHC^{1a} as in systems containing a six-carbon chain between the amino groups and the marked decrease in activity observed after methylation of the inner nitrogen atoms (compounds 23 and 24).

Although removal of the entire (benzylamino)alkyl moieties of BHC to give diamine disulfides 35–38 caused a marked decrease in blockade, probably through a loss of binding ability, these diamines were not completely inactive. The *tert*-butyl derivative of cystamine, 36, was as potent as some tetramine disulfides (compounds 2, 10) that have all the structural requirements for efficient binding. This result seems to indicate that all four protonatable nitrogen atoms may not be essential for activity. To test this possibility, we undertook the syntheses of diamine disulfides 32–34 having oxygen atoms in place of the outer amino groups of BHC. Two of these compounds, 32 and 33, showed a considerable degree of blockade, which seems to deny the necessity for the presence of four anionic centers on the binding area for BHC. Similar results were obtained when substitution of oxygen for nitrogen was carried out in the inner cystamine moiety (compounds 29 and 31) and when the protonable amino groups were changed into nonbasic amide functions (compound 12). In fact, the diamine-diamide 12 is a more potent blocker than the corresponding tetramine 14, a result supportive of a dipolar interaction between the outer portions of tetramine disulfides and their binding areas.

In conclusion, we have found that all compounds containing the central cystamine moiety display an irreversible blockade of the NE contraction of the rat vas deferens. This effect is more apparent in cystamines *N,N'*-disubstituted with 6-aminoheptyl groups, especially when the outer nitrogen atoms bear arylalkyl substituents or are enclosed in a ring. Disubstitution of nitrogen atoms in the cystamine moiety leads to poor blockers. Structural specificity in the outer portion of the tetramine disulfide is low since many types of substituents and even the change of amines into nonbasic groups were observed to be potent α -adrenergic blockers.

Experimental Section

Melting points were determined in a capillary tube on a Büchi apparatus and are uncorrected. Prior to concentration under reduced pressure, all organic extracts were dried over anhydrous $MgSO_4$ powder. TLC and column chromatography were carried out on SiO_2 (silica gel 60, Merck, 63–200 μm) and the compounds were detected in TLC with UV light or iodoplatinate reagent. Molecular distillations were effected by use of a Büchi GKR-50 Kugelrohr apparatus; the temperatures cited are the highest

reached by the oven during distillation. Microanalyses were performed by the Instituto de Química Bio-Orgánica, Barcelona.

General Procedure for Preparation of 3-[[[2-Hydroxy-2-arylethyl]amino]hexyl]-1,3-thiazolidines (45a–h). A solution of the aryloxirane 41a–h (20 mmol) and 3-(6-aminoheptyl)-1,3-thiazolidine (42)^{2a} (20 mmol) in *i*-PrOH (60 mL) was stirred for 22 h at reflux temperature under a N_2 atmosphere. The solvent was removed at reduced pressure; the residue was dissolved in CH_2Cl_2 (100 mL) and extracted with 2 N HCl (4 \times 25 mL). The aqueous layers were basified with 5 N NaOH and extracted with CH_2Cl_2 . The organic extracts were dried and evaporated under reduced pressure to give thiazolidines 45a–h as oils. Purifications were achieved by successive crystallization of the dicarlate derivative.

3-[6-[[1,4-Benzodioxan-2-ylmethyl]amino]hexyl]-1,3-thiazolidines (45i). A solution of (*RS*)-2-(tosylmethyl)-1,4-benzodioxane (43)^{2a} (1.56 mmol) and Et_3N (0.47 g) in *i*-PrOH (20 mL) was stirred at 25 °C under N_2 atmosphere. After a few minutes, a solution of 42 (4.68 mmol) in *i*-PrOH (15 mL) was added and the mixture was stirred at reflux for 24 h. Upon removal of the solvent under reduced pressure, the residue was taken up in CH_2Cl_2 (200 mL) and extracted with 2 N HCl. The aqueous layer was basified with 5 N NaOH and extracted with benzene. The organic extracts were dried and evaporated, and compound 45i was obtained in 86% yield. Enantiomers of this compound were prepared in a similar way, with the corresponding chiral tosylate as the starting material.^{2b}

***N*-[6-(1,3-Thiazolidin-3-yl)hexyl]phenylacetamide (45j) or Benzamides 45k–l.** To a vigorously stirred solution of 42 (2 g, 10.6 mmol) and Et_3N (2.7 g, 26 mmol) in dry benzene (80 mL) was added dropwise a solution of the suitable acid chloride 44a–c (10.6 mmol) in anhydrous benzene. Stirring was continued for 5 h at 25 °C. The reaction mixture was washed with H_2O and extracted with 2 N HCl. The aqueous layer was basified with 2 N NaOH and extracted with benzene. The organic extracts were dried and evaporated, yielding the corresponding compound 45j–l as an oil.

General Procedure for Oxidative Dimerization of Thiazolidines. Preparation of 2–8, 10, 12, and 13. Method A. To a stirred solution of the appropriate thiazolidine 45a–l (20 mmol) in EtOH (200 mL) was added slowly a solution of 0.1 N I_2 /2.5% KI (200 mL) followed by the addition of 2 N HCl (5 mL). The mixture was stirred at 25 °C for 12 h. The solvent was removed at reduced pressure and the residue taken up in 2.5 N NaOH (150 mL). The alkaline solution was washed with CH_2Cl_2 (3 \times 75 mL). The organic extracts were dried and concentrated, giving the corresponding disulfides 2–8, 10, 12, and 13 in high yield. Purifications of these disulfides were achieved by crystallizations of suitable derivatives.

Preparation of *N,N'*-Bis[6-[[2-hydroxy-2-(3,4-dihydroxyphenyl)ethyl]amino]hexyl]cystamine (9). The thiazolidine 45h was transformed by the previous general procedure into the *N,N'*-bis[6-[[2-hydroxy-2-(3,4-bis(methoxymethoxy)phenyl)ethyl]amino]hexyl]cystamine. Through a solution of this cystamine (1.77 g, 2.13 mmol) in dry methanol (40 mL) and methylene chloride (35 mL) was passed a current of dry hydrogen chloride until the solution pH was acidic. The solution was stirred for 20 h at room temperature, the solvent was removed at reduced pressure, and a yellow solid (1.1 g, 60%) was obtained and identified as the tetrahydrochloride of 9.

Preparation of *N,N'*-Bis[6-[(phenylethyl)amino]hexyl]cystamine (14). $LiAlH_4$ (0.58 g, 15.3 mmol) was slowly added to a solution of 12 (3 g, 5 mmol) in dioxane (80 mL), and the resulting mixture was stirred at reflux temperature for 4 h. After cooling, excess $LiAlH_4$ was destroyed with H_2O , and the resulting mixture was basified with 2 N NaOH. The organic layer was separated, dried, and evaporated to give compound 14 as an oil (2.01 g, 95%).

Preparation of *N,N'*-Bis[6-[(*o*-aminobenzyl)amino]hexyl]cystamine (15). The thiazolidine 45k was transformed by the previous general procedure into the *N,N'*-bis[6-[(*o*-nitrobenzyl)amino]hexyl]cystamine. A solution of this cystamine (4.39 g, 6 mmol) in dry dioxane (150 mL) was slowly added to a suspension of $LiAlH_4$ (3.59 g, 90 mmol) in dry dioxane (70 mL). The mixture was stirred at reflux temperature for 5 h. A nitrogen atmosphere was maintained in a system during the entire process.

The solution was cooled and H_2O (3.6 mL), dioxane (3.6 mL), 2 N NaOH (7.8 mL), and H_2O (10 mL) were added in the order indicated. The suspension that formed was filtered, the solid was washed several times with CHCl_3 , all the liquid fractions were combined and dried, and the solvent was removed under vacuum to give 15 as an oil (1.6 g, 43%).

Preparation of N,N' -Bis[6-[(*o*-aminobenzyl)amino]hexyl]- N,N' -dimethylcystamine (16). LiAlH_4 (4.46 g, 117 mmol) was slowly added to a solution of 451 (4.42 g, 13 mmol) in dry dioxane (200 mL) and the mixture was stirred at reflux temperature under N_2 atmosphere for 5 h. The solution was cooled, and dioxane (4.5 mL), H_2O (4.5 mL), 2 N NaOH (7 mL), and H_2O (13 mL) were added in the order indicated. The suspension that formed was filtered, the solid was washed several times with CHCl_3 , and the combined organic layers were dried and evaporated to give an oil (3 g). This oil was dissolved in EtOH (50 mL) and over this ethanolic solution 0.1 N aqueous I_2 was added until no decoloration was observed. The solution was stirred for 1 h and evaporated and the residue was dissolved in H_2O . The aqueous solution was basified with 2 N NaOH and extracted with CH_2Cl_2 . The organic layer was dried and evaporated to give the disulfide 16 (2.16 g, 56%).

General Procedure for Preparation of Cystamines 17–20. Method B. A solution of N,N' -bis(6-aminohexyl)cystamine (46)^{1a} (10 mmol) in dry benzene (140 mL) was added dropwise to a solution of the suitable heterocyclic aldehyde 47a–e (20 mmol) in dry benzene. The mixture was stirred at reflux temperature for 5 h. The solvent was removed at reduced pressure, and the residue was dissolved in MeOH (160 mL). The solution was cooled to 0 °C, NaBH_4 (24 mmol) was added, and the mixture was stirred at room temperature for 1 h. Concentrated HCl was added until acid pH was reached. MeOH was removed at reduced pressure and the residue washed with CH_2Cl_2 . The aqueous layer was basified with 2 N NaOH and extracted with CH_2Cl_2 . Organic extracts were dried and evaporated to give 17–20, which were purified by crystallization of a suitable derivative.

General Procedure for Preparation of Amino Amides 48a–d. A solution of 6-phthalimidohexanoyl chloride (140 mmol) in dry CH_2Cl_2 (250 mL) was slowly added to a solution of the heterocyclic amine (piperidine, pyrrolidine, tetrahydroisquinoline, or indoline) (140 mmol) and Et_3N (420 mmol) in dry CH_2Cl_2 (200 mL), and the resulting mixture was stirred at 25 °C for 5 h. The organic solution was washed with H_2O , 10% HCl, and 1 N NaOH. Drying and evaporation gave good yields of phthalimido amides,

which were used without further purification for the next reaction. To a solution of the phthalimido amide product (118 mmol) in EtOH (620 mL) was added 80% aqueous hydrazine (118 mmol) and the mixture was stirred at reflux temperature for 3 h. The solvent was evaporated, the residue dissolved in 5 N NaOH, and the product extracted with CHCl_3 . The organic layer was dried and evaporated to give the corresponding heterocyclic amino amide 48a–d.

Preparation of Diamides 48e,f. A mixture of Ac_2O (27.2 g, 260 mmol) and formic acid (10.6 mL, 260 mmol) was stirred for 4 h at 55 °C. After cooling, a solution of the appropriate amide 48a,b (110 mmol) in dry THF (200 mL) was added and the resulting solution was stirred at reflux temperature for 20 h. The organic solution was washed several times with an aqueous solution of Na_2CO_3 , dried, and evaporated to give 48e,f.

General Procedure for Preparation of Cystamines 21–24, 28, and 29 (Method C). A solution of the appropriate amino amide (48a–d) or diamide (48e,f) (25 mmol) in dry THF was added to a suspension of LiAlH_4 (126 mmol) in dry THF (250 mL) and the resulting mixture was stirred at reflux temperature for 5 h. After the mixture cooled to 25 °C, excess LiAlH_4 was destroyed with H_2O and the mixture was filtered. The organic layer was dried and evaporated to give the corresponding N -(6-aminohexyl) heterocycle as an oil, which was purified by vacuum distillation.

A solution of thirane (11 mmol) in dry benzene (15 mL) was slowly added to a solution of N -(6-aminohexyl) heterocycle (11 mmol) in benzene (25 mL), and the mixture was stirred for 14 h at reflux temperature. The solvent was evaporated, the resulting oil was dissolved in EtOH (100 mL), and aqueous 0.1 N I_2 was added over the solution until no decoloration was observed. The solvent was evaporated, the resulting solid was dissolved in 2 N NaOH and extracted with CH_2Cl_2 , and the resulting organic layer

was dried and evaporated. Distillation of the residue allowed recovery of a quantity of the starting diamine. Purification of the product residue by crystallization of a suitable solid derivative allowed us to obtain cystamines 21–24, 28, 29.

Preparation of Cystamines 28–27. A 40% aqueous solution of HCHO (10 mmol) was added to a solution of the suitable compound 15, 16, or 49^{1c} (3 mmol) and KOH (0.23 g, 4 mmol) in MeOH (50 mL). The mixture was stirred at reflux temperature for 1 h, MeOH was removed, and H_2O was added to the resulting residue. The aqueous layer was extracted with CHCl_3 , and the organic solvent was dried and evaporated to give the cystamines 25–27, respectively.

Preparation of N,N' -Bis(6-amino-7-phenylheptyl)cystamine (11). Method D. (a) 3-(7-Phenyl-6-oxoheptyl)-1,3-thiazolidine (50). A 1.26 M solution of $n\text{-BuLi}$ in hexane (18.2 mL) was slowly added to a solution of 2-benzyl-1,3-dithiane⁴ (6.2 g, 20 mmol) in THF (80 mL) at –35 °C and stirring was continued for 35 min. The resulting solution was added to a –10 °C solution of 1,5-dibromopentane (14.5 g, 62 mmol) in THF (100 mL). The mixture was stirred at this temperature for 30 min, warmed to 2–3 °C, and stirred for 21 h. A N_2 atmosphere was maintained in the system throughout the entire process. Reaction workup required that H_2O (6 mL) be added and the solvent separated, and the residue was dissolved in H_2O , acidified with 2 N HCl, and extracted with CHCl_3 . The organic layer was successively washed with an aqueous solution of 5% Na_2SO_3 , 5% KOH, and H_2O . Drying and evaporation yielded an oil, which when distilled at 70 °C (0.09 mmHg) yielded 1,5-dibromopentane. The remaining residue was the desired 2-benzyl-2-(5-bromopentyl)-1,3-dithiane (82%). A suspension of this dithiane (4.4 g, 12 mmol), HgCl_2 (14.3 g, 52 mmol), and HgO (4.2 g, 19 mmol) in MeOH (300 mL) and H_2O (18 mL) was stirred under a N_2 atmosphere at reflux temperature for 4 h. The solution was cooled and filtered and the solid washed with CH_2Cl_2 . The organic layers were evaporated, and the residue was dissolved in H_2O and extracted with CH_2Cl_2 . This solution was successively washed with saturated aqueous NH_4Cl solution, 10% HCl, 10% NaOH, and H_2O . The organic layer upon drying and evaporation gave 7-bromo-1-phenyl-2-heptanone (86%). A suspension of this bromo ketone (3.8 g, 14 mmol), 1,3-thiazolidine (1.4 g, 16 mmol), and anhydrous K_2CO_3 (1.7 g, 16 mmol) in *i*-PrOH (150 mL) was stirred at reflux temperature for 23 h under N_2 atmosphere. The mixture was filtered and the solution evaporated to dryness. The residue was dissolved in CHCl_3 and extracted with 10% HCl. The aqueous layer was basified with 2 N NaOH and extracted with CHCl_3 . The organic layer when dried and evaporated gave 50 as an oil.

(b) N,N' -Bis(7-phenyl-6-oxoheptyl)cystamine (51). The oil 50 was dissolved in EtOH and to this solution was added an aqueous 0.1 N I_2 solution (83 mL). After the mixture was stirred for 2 h, the solvent was eliminated, and the residue was basified with 2 N NaOH and extracted with CHCl_3 . The organic layer was dried and evaporated to give N,N' -bis(7-phenyl-6-oxoheptyl)cystamine (51) (56%).

(c) **Preparation of Cystamine 11.** A solution of diketone 51 (2.5 g, 4.8 mmol) in pyridine/EtOH (2:1) (46 mL) and hydroxylamine hydrochloride (1 g, 14.4 mmol) was stirred at reflux temperature for 5 h. The solution was evaporated to dryness, and the residue was dissolved in 1 N NaOH and extracted with CHCl_3 . The organic layer was washed with H_2O , dried, and evaporated to give the intermediate oxime. The oxime was dissolved in THF (75 mL) and dioxane (25 mL) and LiAlH_4 (0.34 g, 8.9 mmol) was added. The mixture was stirred at reflux temperature for 1 h under N_2 atmosphere. The suspension was cooled, and 2 mL of H_2O and 2 mL of 2 N NaOH were successively added. It was filtered and the residue dissolved in CHCl_3 . The combined organic layers were washed with H_2O , dried, and evaporated to give the cystamine 11 (20%).

Preparation of Bis[2-[(*o*-[(arylmethyl)amino]hexyl]oxy]ethyl] Disulfides 30 and 31. Method E. (a) 11-Phenyl-7-oxa-10-thiaundecylamine (53). Under N_2 atmosphere, a solution of 2-(benzylthio)ethanol⁵ (52) (8.4 g, 50 mmol) in THF (35 mL) and DMF (15 mL) was slowly added to a suspension of NaH (1.4 g, 59 mmol) in THF (35 mL) and DMF (15 mL). The resulting mixture was stirred for 5 min after which a solution of 1,5-dibromopentane (22.9 g, 100 mmol) in THF (35 mL) and DMF (15 mL) was slowly added. The reaction mixture

was warmed to reflux temperature and stirred for 1 h. After this time H_2O and Et_2O were added. The organic layer was dried and evaporated to give 30.8 g of an oil. Distillation of the oil gave 12.5 g of 1,5-dibromopentane (85 °C, 0.07 mmHg) and a residue, which was purified by column chromatography; on elution with CHCl_3 , 6.2 g of 10-bromo-1-phenyl-5-oxa-2-thiadecane was obtained. A solution of KCN (1.5 g, 23 mmol) in H_2O (63 mL) was added to a solution of the above alkyl bromide (6.2 g, 20 mmol) in EtOH (145 mL) and the mixture was stirred at reflux temperature for 4 h. After this time, the EtOH was removed, H_2O was added, and the compound was extracted with Et_2O . The organic layer was dried and evaporated to give 4.9 g of 11-phenyl-7-oxa-10-thiaundecanenitrile (38%). A solution of this nitrile (4.9 g, 18 mmol) in dioxane (50 mL) was added to a suspension of LiAlH_4 (2.1 g, 56 mmol) in dioxane (100 mL) and the resulting mixture was stirred at reflux temperature for 3 h. After cooling, 3 mL of H_2O , 3 mL of dioxane, 4 mL of 2 N NaOH, and 9 mL of H_2O were successively added. The mixture was filtered and the filtrate evaporated to dryness. The residue was dissolved in CHCl_3 and extracted with 10% HCl. The aqueous layer was basified with 2 N NaOH and washed with CHCl_3 . The chloroform extracts were dried and evaporated to give 3.29 g of an oil. This oil was distilled and the fraction collected between 175 and 200 °C (0.05 mmHg) yielded 2.26 g of 53 (44%).

(b) Bis[2-[(6-aminohexyl)oxy]ethyl] Disulfide (54). Under N_2 atmosphere, small pieces of Na (4.5 g, 195 mmol) were added to a solution of 53 (0.5 g, 1.8 mmol) in EtOH (56 mL). The reaction was strongly exothermic. When spontaneous boiling ceased, the solution was maintained at reflux temperature until all of the Na had dissolved (40 min). The solution was cooled and solid CO_2 was added. The resulting white paste was diluted with EtOH and CH_2Cl_2 and filtered, and the solid was washed with CH_2Cl_2 . The organic layers were evaporated to dryness, and the residue was dissolved in EtOH . Over this ethanolic solution was added 0.1 N aqueous I_2 until no decoloration was observed, and the mixture was stirred for 1 h. The solution was evaporated, and the residue was dissolved in H_2O , basified with 2 N NaOH, and extracted with CH_2Cl_2 . The organic layer was dried and evaporated to give the disulfide 54 (0.3 g, 90%).

(c) Bis[2-[[6-[(arylmethyl)amino]hexyl]oxy]ethyl] Disulfides 30 and 31. A solution of disulfides 54 (0.3 g, 0.9 mmol) and the suitable aldehyde (benzaldehyde or *o*-methoxybenzaldehyde) (1.9 mmol) in dry benzene (50 mL) was stirred at reflux temperature for 15 h in a system equipped with a Dean-Stark trap. After this time the solvent was evaporated and the residue dissolved in MeOH . To this methanolic solution was added NaBH_4 (0.08 g, 2.1 mmol) and the mixture was stirred at reflux temperature for 30 min. The mixture was cooled, acidified with 2 N HCl, and washed with Et_2O . The aqueous solution was basified with 2 N NaOH and extracted with CHCl_3 . The chloroform extracts were dried and evaporated to give the expected disulfides 30 or 31.

General Procedure for Preparation of N,N' -Bis[(arylmethoxy)hexyl]cystamines 32-38. Method F. (a) [(Arylmethoxy)hexyl]amines 56a-c. A solution of the suitable alcohol 55a-c (80 mmol) in anhydrous THF (20 mL) was slowly added to a suspension of NaH (100 mmol) and 1,6-dibromohexane (200

mmol) in THF (60 mL). The resulting mixture was stirred at reflux temperature for 30 min, then poured into H_2O , and extracted with Et_2O . The organic layer was washed with H_2O , dried, and evaporated to give an oil. Upon distillation, excess 1,6-dibromopentane and a residue identified as the corresponding bromo ether were isolated. The bromo ether and potassium phthalimide (50 mmol) in DMF (50 mL) were heated at 100 °C for 5 h. After this time the mixture was poured into H_2O and extracted with Et_2O . The organic layer was washed with H_2O , dried, and evaporated to give the corresponding phthalimide. A solution of this phthalimide (60 mmol) and 80% hydrazine hydrate (104 mmol) in EtOH (300 mL) was heated for 3 h at reflux temperature. The solvent was removed, the residue was dissolved in 5 N NaOH, and the desired product was extracted with CHCl_3 . The CHCl_3 extracts were dried and evaporated to give the corresponding amines 56a-c.

(b) Cystamines 32-38. A solution of the suitable amine (56a-c, ethylamine, *tert*-butylamine, benzylamine, or piperidine) (75 mmol) in benzene (20 mL) was stirred at reflux temperature for 1 h in a system equipped with a Dean-Stark trap.

The solution was cooled at 0 °C and a thiirane (63 mmol) in benzene (20 mL) solution was slowly added. The resulting mixture was refluxed for 3 h. The solvent was evaporated, the residue was dissolved in H_2O , and 2 N HCl was added until a pH of 8-9 was reached. A solution of $\text{K}_3\text{Fe}(\text{CN})_6$ (8 mmol) in H_2O (25 mL) was added. The mixture was allowed to stand, and after a 30-min period, NaOH pellets (6 g, 150 mmol) were added. A sufficient amount of NaCl was next added to saturate the solution. The aqueous solution was washed with CH_2Cl_2 , and the organic extracts were dried and evaporated to give disulfides 32-38. These compounds were purified by crystallization of a solid derivative.

Pharmacology. The following protocol^{1a} was applied for the relative potencies listed in Table I. Male rats weighing 200-250 g were killed by a sharp blow on the head and both vasa deferentia were isolated. These were mounted individually in organ baths of 30-mL vol containing Krebs bicarbonate buffer (113 mmol of NaCl, 4.7 mmol of KCl, 2.4 mmol of CaCl_2 , 1.2 mmol of MgSO_4 , 1.2 mmol of KH_2PO_4 , 25 mmol of NaHCO_3 , and 11.5 mmol of dextrose). The medium was maintained at 32 °C while being aerated with 95% O_2 -5% CO_2 . The loading tension was 0.50 g, and the contractions were recorded by means of force transducers connected to a Omni-Scribe recorder. The tissues were allowed to equilibrate for 1 h, and the medium was changed prior to addition of the antagonists. After a 30-min incubation period, the bath was drained and the tissues were washed with the bath solution for 30 min. Cumulative concentration-response curves for NE were constructed after treatment with each antagonist. The decrease in maximum response was expressed in percent of the control value. The percent blockade for each compound is expressed as the mean \pm SEM of five separate experiments.

Acknowledgment. We thank Dr. F. López-Calabrera, for his work and encouragement in the initial step, A. Morón, E. Naudó, E. Bardají, and L. Bennasar for their assistance with the synthetic work, and also the Department of Pharmacology, Autònoma University of Barcelona, for the use of their facilities.

Modifications of Primaquine as Antimalarials. 4. 5-Alkoxy Derivatives of Primaquine

Eugene H. Chen, Keiichi Tanabe, Andrew J. Saggiomo, and Edward A. Nodiff*

Franklin Research Center, Division of Arvin/Calspan, Philadelphia, Pennsylvania 19103. Received October 3, 1986

Thirty-two 5-alkoxyprimaquines have been synthesized and evaluated as blood schizonticides (*Plasmodium berghei*, mouse) and tissue schizonticides (*Plasmodium cynomolgi*, monkey). Several of these compounds were extremely active in both screens. Such a broad spectrum of antimalarial efficacy offers the possibility of a single drug that could cure the various relapsing and nonrelapsing malarias.

The major forms of human malaria are caused by the parasites *Plasmodium falciparum* and *Plasmodium vivax*.

Elimination of the blood form of falciparum malaria clears the body of this disease. Vivax malaria, however, gives rise